

Address the appearance of aging skin with the power of Vitamin C.

Vitamin C is an important antioxidant¹

- Calms skin²⁻⁴
- Brightens skin^{5,6}
- Retains moisture⁷
- Stabilizes Vitamin E⁸
- Reduces the appearance of fine lines and wrinkles^{9,10}

	Calms skin	Brightens skin	Retains moisture	Stabilizes Vitamin E	Reduces the look of fine lines and wrinkles
L-ascorbic acid (Vitamin C) ²⁻¹⁰	✓	✓	✓	✓	✓
Idebenone ¹¹			✓		✓
Coffee berry ¹²		✓			
Kojic acid ¹³		✓			
Green tea ¹²	✓				
Emblica (Indian gooseberry) ^{14,15}		✓			✓
Niacinamide ¹²	✓	✓			

References: **1.** Lewis AB, Regan K. Dermatological procedure enhancement products—a beneficial means to combat aging skin. *US Dermatol Review.* 2006;1-5. **2.** Alster TS, West TB. Effect of topical vitamin C on postoperative CO2 laser resurfacing erythema. **3.** Basketter DA, et al. Influence of vitamin C on the elicitation of allergic contact dermatitis to p-phenylenediamine. *Contact Dermatitis.* 2016 Jun;74(6):368-72. **4.** Coenraads PJ, et al. The role of the antioxidant ascorbic acid in the elicitation of contact allergic reactions to p-phenylenediamine. *Contact Dermatitis.* 2016 May;74(5):267-72. **5.** Kameyama K, Sakai C, Kondoh S, et al. Inhibitory effect of magnesium L-ascorbyl-2-phosphate (VC-PMG) on melanogenesis in vitro and in vivo. *J Am Acad Dermatol.* 1996;34(1):29-33. **6.** Espinal-Perez LE, Moncada B, Castanedo-Cazares JP. A double-blind randomized trial of 5% ascorbic acid vs. 4% hydroquinone in melasma. *Int J Dermatol.* 2004 Aug;43(8):604-7. **7.** Campos PM, Gonçalves GM, Gaspar LR. In vitro antioxidant activity and in vivo efficacy of topical formulations containing vitamin C and its derivatives studied by non-invasive methods. *Skin Res Technol.* 2008;14(3):376-380. **8.** Burgess C. Topical vitamins. *J Drugs Dermatol.* 2008;7(7 suppl):52-56. **9.** Traikovich S. Use of Topical Ascorbic Acid and Its Effects on Photodamaged Skin Topography. *Arch Otolaryngol Head Neck Surg.* 1999;125(10):1091-1098. **10.** Farris PK. Topical vitamin C: a useful agent for treating photoaging and other dermatologic conditions. *Dermatol Surg.* 2005;31(7, pt 2):814-818. **11.** McDaniel DH, et al. Clinical efficacy assessment in photodamaged skin of 0.5% and 1.0% idebenone. *J Cosmet Dermatol.* 2005 Sep;4(3):167-73. **12.** Bogdan Allemann I, Baumann L. Antioxidants used in skin care formulations. *Skin Therapy Lett.* 2008;13(7):5-9. **13.** Halder RM, Richards GM. Topical agents used in the management of hyperpigmentation. *Skin Therapy Lett.* 2004;9(6):1-3. **14.** Skin firming properties reported in Indian gooseberry. *Cosmetics and Toiletries Science Applied website.* <http://www.cosmeticsandtoiletries.com/formulating/function/active/38024659.html>. Accessed July 9, 2015. **15.** Draelos ZD, Yatskayer M, Bhushan P, et al. Evaluation of a kojic acid, emblica extract, and glycolic acid formulation compared with hydroquinone 4% for skin lightening. *Cutis.* 2010;86(3):153-158.

All products/brand names, whether designated by notice or not (®/™), are trademarks of Obagi Cosmeceuticals LLC and/or its affiliates, except for those that are trademarks of their respective owners, which are used by permission. Distributed by Obagi Cosmeceuticals LLC.

©2019 Obagi Cosmeceuticals LLC. All rights reserved. www.obagi.com 03/19 PRF.0011.USA.19.r0.

